



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

STATISTICAL REPORTS OF MORBIDITY AND MORTALITY, STATES OF THE UNITED STATES (Untabulated).

FLORIDA.—Reports received from the State board of health show as follows: Week ended April 29, 1912. Diphtheria was reported present at 2 places with 4 cases; malaria at 5 places with 53 cases; tuberculosis at 6 places with 14 cases; typhoid fever at 5 places with 7 cases.

Smallpox.—During the month of April, 1912, smallpox was reported in 7 counties with 82 cases.

Week ended May 6, 1912. Diphtheria present at 1 place (West Palm Beach) with 1 case; malaria at 5 places with 38 cases; smallpox 29 cases in 6 counties; tuberculosis at 8 places with 9 cases; typhoid fever at 2 places with 3 cases.

Week ended May 12, 1912. Diphtheria present at 5 places with 6 cases; malaria at 7 places with 48 cases; tuberculosis at 6 places with 7 cases; typhoid fever at 6 places with 8 cases.

IOWA.—Month of March, 1912. Population, 2,224,771. Total number of deaths from all causes, 2,092, including diphtheria 8, measles 3, scarlet fever 18, tuberculosis 63, typhoid fever 23.

MINNESOTA.—Month of January, 1912. Population, 2,075,708. Total number of deaths from all causes, 1,745, including diphtheria 19, measles 2, scarlet fever 18, tuberculosis 205, typhoid fever 23.

Month of February, 1912. Total number of deaths from all causes, 1,584, including diphtheria 21, measles 1, scarlet fever 10, smallpox 3, tuberculosis 198, typhoid fever 16.

UTAH.—Month of April, 1912. Population, 373,351. Total number of deaths from all causes, 320, including diphtheria 1, measles 1, scarlet fever 4, tuberculosis 8, typhoid fever 1. Cases reported: Diphtheria 24, measles 88, scarlet fever 61, smallpox 267, tuberculosis 7 (incomplete), typhoid fever 10.